arXiv:1404.7466v4 [math.AG] 8 Mar 2015

ON FINITE REGULAR AND HOLOMORPHIC MAPPINGS

ZBIGNIEW JELONEK

Abstract. Let X, Y be smooth algebraic varieties of the same dimension. Let $f, g : X \longrightarrow Y$ be finite regular mappings. We say that f, g are equivalent if there exists a regular automorphism $\Phi \in \mathrm{Aut}(X)$ such that $f = g \circ \Phi$. Of course if f, g are equivalent, then they have the same discriminants (i.e., the same set of critical values) and the same geometric degree. We show that conversely, for every hypersurface $V \subset Y$ and every $k \in \mathbb{N}$, there are only a finite number of non-equivalent finite regular mappings $f : X \to Y$ such that the discriminant $D(f)$ equals V and $\mu(f) = k$. As one of applications we show that if $f : X \to Y$ is a finite mapping of topological degree two, then there exists a regular automorphism $\Phi : X \to X$ which acts transitively on the fibers of f and $Y = X/G$, where $G = \{id, \Phi\}$ and f is equivalent to the canonical projection $X \to X/G$.

We prove the same statement in the local (and sometimes global) holomorphic situation. In particular we show that if $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ is a proper and holomorphic mapping of topological degree two, then there exist biholomorphisms $\Psi, \Phi : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ such that $\Psi \circ f \circ \Phi(x_1, x_2, \ldots, x_n) = (x_1^2, x_2, \ldots, x_n)$. Moreover, for every proper holomorphic mapping $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ with smooth discriminant there exist biholomorphisms $\Psi, \Phi : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ such that $\Psi \circ f \circ \Phi(x_1, x_2, \ldots, x_n) = (x_1^k, x_2, \ldots, x_n)$, where $k = \mu(f)$.

1. Introduction

1.1. **Global case.** Let X, Y be smooth algebraic (resp. holomorphic) varieties of the same dimension. We say that a regular (resp. holomorphic) mapping $f : X \to Y$ is *finite*, if it is proper and it has finite fibers. If X, Y are affine (resp. Stein) then every proper mapping is finite. Let $f, g : X \to Y$ be finite regular (resp. holomorphic) mappings. By the *discriminant* $D(f)$ of f we mean the set of critical values of f. Of course $D(f)$ is a hypersurface in Y. For every $y \in U = Y \setminus D(f)$ the fiber $f^{-1}(y)$ has exactly $\mu(f)$ points, where $\mu(f)$ is the topological degree of f. In the case $X = Y = \mathbb{C}^2$, S. Lamy [Lam] (see also [B-C] and [B-C-1]) proved the following theorem:

Theorem 1.1. *Let $f : \mathbb{C}^2 \to \mathbb{C}^2$ be a finite polynomial map of topological degree 2. Then there exist polynomial automorphisms Φ_1, Φ_2 such that $f = \Phi_2 \circ g \circ \Phi_1$, where $g(x, y) = (x, y^2)$.*

This theorem, although interesting, is very special. We explain why it is so special and we generalize it in a few directions (see Theorem 1.15, Theorem 1.16 and Corollary 1.7). In fact to describe a finite regular mapping it is not enough to use only the geometric degree. We show that in principle a finite regular mapping is determined by its discriminant and the geometric degree. The same is true in a local (and sometimes in a global) holomorphic case.

Definition 1.2. *Let X, Y be smooth algebraic (resp. holomorphic) manifolds of the same dimension. Let $f, g : X \to Y$ be finite regular (resp. holomorphic) mappings. We say*

Date: March 10, 2015.

1991 *Mathematics Subject Classification.* 14 A 10, 14 R 10, 51 M 99.

The author was partially supported by the NCN grant 2014-2017.

that f, g are equivalent if there exists a regular automorphism (resp. a biholomorphism) $\Phi : X \to X$ such that $f = g \circ \Phi$. Moreover, we say that f is weakly equivalent to g if there exist regular automorphisms (resp. biholomorphisms) $\Phi \in Aut(X), \Psi \in Aut(Y)$ such that $g = \Psi \circ f \circ \Phi$.

Of course if f, g are equivalent, then they have the same discriminant and the same geometric degree. Our first main result is the following:

Theorem 1.3. *Let X, Y be smooth algebraic varieties of the same dimension. Let $V \subset Y$ be a hypersurface and k be a natural number. There are only a finite number of non-equivalent finite regular mappings $f : X \to Y$ such that $D(f) = V$ and $\mu(f) = k$.*

Remark 1.4. The same result is true in the holomorphic category, if we additionally assume that the fundamental group of the space $Y \setminus V$ is finitely generated.

Note that in general there are two or more non-equivalent proper mappings with the same discriminant and with the same geometric degree. For example for $X = Y = \mathbb{C}^2$ there are exactly three non-equivalent proper mappings $f_1(x, y) = (x^4, y^4)$, $f_2(x, y) = (x^8, y^2)$ and $f_3(x, y) = (x^2, y^8)$ with $D(f) = \{(x, y) \in \mathbb{C}^2 : xy = 0\}$ and $\mu(f) = 16$ (see Corollary 1.7). However sometimes there is only one such a map:

Corollary 1.5. *Let $V = \{x \in \mathbb{C}^n : x_1 = 0\}$. Every proper polynomial (resp. holomorphic) mapping $f : \mathbb{C}^n \to \mathbb{C}^n$ with $D(f) = V$ and $\mu(f) = k$ is equivalent to the mapping*

$$g(x_1, x_2, \ldots, x_n) = (x_1^k, x_2, \ldots, x_n).$$

This result has the following more general counterpart:

Corollary 1.6. *Let $V = \{x \in \mathbb{C}^n : x_1 = 0\}$ and let X be an n-dimensional smooth affine (resp. Stein) variety. Then any two proper regular (resp. holomorphic) mappings $f, g : X \to \mathbb{C}^n$ with $D(f) = V$ and $\mu(f) = \mu(g) = k$ are equivalent. Moreover, there is then a regular automorphism (resp. a biholomorphism) $\sigma : X \to X$ of order k such that $f \circ \sigma = f$ (i.e., σ acts transitively on the fibers of f).*

We have here one of possible generalizations of Theorem 1.1 (for other generalizations see Theorems 1.15 and 1.16):

Corollary 1.7. *Let X, Y be smooth algebraic (resp. holomorphic) manifolds of dimension n. Let $f : X \to Y$ be a finite regular (resp. holomorphic) mapping with $\mu(f) = 2$. Then there is an automorphism (resp. a biholomorphism) $\sigma : X \to X$ of order two such that $f \circ \sigma = f$ (i.e., σ acts transitively on fibers of f). The set of fixed points of σ coincides with the set of critical points of f.*

As a nice application of Corollary 1.7 we have:

Corollary 1.8. *If $f : X \to Z$ (where X is smooth and Z is normal) is a finite surjective morphism (resp. holomorphic mapping) of topological degree two, then there is a subgroup $G \subset Aut(X)$ of order two, such that*

1) $Z \cong X/G$,

2) f is equivalent to the projection $\pi : X \to X/G$.

We also have:

Corollary 1.9. *Let $K_n = \{x \in \mathbb{C}^n : \prod_{i=1}^n x_i = 0\}$. Every proper polynomial (resp. holomorphic) mapping $f : \mathbb{C}^n \to \mathbb{C}^n$ with $D(f) = V$ and $\mu(f) = k$ is equivalent to one of the mappings*

$$f_{d_1,\ldots,d_n} : \mathbb{C}^n \ni (x_1, \ldots, x_n) \mapsto (x_1^{d_1}, \ldots, x_n^{d_n}) \in \mathbb{C}^n,$$

where $\prod_{i=1}^n d_i = k$ *and all* $d_i > 1$. *In particular if* k *is a prime number and* $n > 1$, *then there is no proper polynomial (resp. holomorphic) mappings with* $D(f) = K_n$ *and* $\mu(f) = k$.

Our method can also sometimes be applied to arbitrary regular mappings with finite fibers (i.e., quasi-finite mappings). Let us recall that if $f : \mathbb{C}^n \to \mathbb{C}^n$ is a generically-finite regular mapping, then $B(f) = \{x \in \mathbb{C}^n : \#f^{-1}(x) \neq \mu(f)\}$ is called the bifurcation set of f. It is always a hypersurface (see e.g. [J-K]). We have:

Corollary 1.10. *Let* $K_n = \{x \in \mathbb{C}^n : \prod_{i=1}^n x_i = 0\}$. *Every quasi-finite regular mapping* $f : \mathbb{C}^n \to \mathbb{C}^n$ *with* $B(f) \subset K_n$ *is proper.*

If we use the concept of weak equivalence we have:

Corollary 1.11. *a) Let* $f : \mathbb{C} \to \mathbb{C}$ *be a polynomial. If* $D(f)$ *is a point, then* f *is weakly equivalent to the mapping* $g(x) = x^k$, *where* $k = \mu(f)$. *Also there are only a finite number of non weakly equivalent polynomials* f *of degree* k *which have exactly two critical points.*

b) Let $f : \mathbb{C}^2 \to \mathbb{C}^2$ *be a proper polynomial mapping. If* $D(f)$ *is isomorphic to* $\mathbb{A}^1(\mathbb{C})$, *then* f *is weakly equivalent to the mapping* $g(x, y) = (x^k, y)$, *where* $k = \mu(f)$.

c) Let $f : \mathbb{C}^n \to \mathbb{C}^n$ *be a proper polynomial mapping. If* $D(f)$ *is isomorphic to* $K_n = \{x \in \mathbb{C}^n : \prod_{i=1}^n x_i = 0\}$, *then* f *is weakly equivalent to one of the mappings*

$$f_{d_1,\dots,d_n} : \mathbb{C}^n \ni (x_1, \dots, x_n) \mapsto (x_1^{d_1}, \dots, x_n^{d_n}) \in \mathbb{C}^n,$$

where $\prod_{i=1}^n d_i = \mu(f)$ *and all* $d_i > 1$.

1.2. **Local case.** It is very interesting that our method works in the local holomorphic setting as well. In particular we will prove:

Theorem 1.12. *Let* $(V, 0) \subset (\mathbb{C}^n, 0)$ *be a germ of analytic hypersurface and let* k *be a natural number. There are only a finite number of non-equivalent proper holomorphic mappings* $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ *such that* $D(f) = V$ *and* $\mu(f) = k$.

Corollary 1.13. *Let* $V = \{x \in \mathbb{C}^n : x_1 = 0\}$. *Every proper holomorphic mapping* $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ *with* $D(f) = (V, 0)$ *and* $\mu(f) = k$ *is equivalent to the mapping*

$$g(x_1, x_2, \dots, x_n) = (x_1^k, x_2, \dots, x_n).$$

In particular every proper holomorphic mapping $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ *with smooth discriminant is weakly equivalent to the mapping*

$$g(x_1, x_2, \dots, x_n) = (x_1^k, x_2, \dots, x_n),$$

where $k = \mu(f)$.

Corollary 1.14. *Let* $K_n = \{x \in \mathbb{C}^n : \prod_{i=1}^n x_i = 0\}$. *Every proper holomorphic mapping* $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ *with* $D(f) = (K_n, 0)$ *and* $\mu(f) = k$ *is equivalent to one of the mappings*

$$f_{d_1,\dots,d_n} : \mathbb{C}^n \ni (x_1, \dots, x_n) \mapsto (x_1^{d_1}, \dots, x_n^{d_n}) \in \mathbb{C}^n,$$

where $\prod_{i=1}^n d_i = k$ *and all* $d_i > 1$. *In particular every proper holomorphic mapping* $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ *with* $D(f)$ *biholomorphic to* $(K_n, 0)$ *and* $\mu(f) = k$ *is weakly equivalent to one of these mappings.*

Hence if k *is a prime number and* $n > 1$, *then there is no proper holomorphic mapping* $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ *with* $D(f)$ *biholomorphic to* $(K_n, 0)$ *and* $\mu(f) = k$.

We also have the following interesting:

Theorem 1.15. *Let $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ be a proper holomorphic mapping with $\mu(f) = 2$. Then f is weakly equivalent to the mapping*

$$g : (\mathbb{C}^n, 0) \ni (x_1, x_2, \ldots, x_n) \mapsto (x_1^2, x_2, \ldots, x_n) \in (\mathbb{C}^n, 0).$$

In particular the discriminant of f is smooth.

One can also prove an algebraic counterpart of Theorem 1.15. To do this we need:

Linearization Conjecture. *Let $\Phi : \mathbb{C}^n \to \mathbb{C}^n$ be a polynomial automorphism of order two. Then there exists an automorphism $\Psi : \mathbb{C}^n \to \mathbb{C}^n$ such that*

$$\Psi^{-1} \circ \Phi \circ \Psi \in GL(n).$$

This Conjecture is true in dimension 2 (and $n = 1$ of course) - see [Kam], but unfortunately it is completely open in higher dimensions. We have the following generalization of the Lamy Theorem:

Theorem 1.16. *Let $f : \mathbb{C}^n \to \mathbb{C}^n$ be a proper polynomial mapping with $\mu(f) = 2$. If the Linearization Conjecture is true, then f is weakly equivalent to the mapping*

$$g : \mathbb{C}^n \ni (x_1, x_2, \ldots, x_n) \mapsto (x_1^2, x_2, \ldots, x_n) \in \mathbb{C}^n.$$

Sections 2 and 3 are devoted to the proofs of Theorem 1.3 and Corollaries 1.4-1.11. In Section 4 we prove Theorem 1.12 and Corollaries 1.13-1.14. In section 5 we prove Theorems 1.15 and 1.16.

Remark 1.17. Our methods also work if we do not assume that Y is smooth and we define the discriminant of a finite mapping $f : X \to Y$ as $D(f) = \{y \in Y : y \in \text{Sing}(Y) \text{ or } \#f^{-1}(y) < \mu(f)\}$.

2. Proof of Theorem 1.3

Let $V \subset X$ be a hypersurface. It is well known that the fundamental group of a smooth algebraic variety is finitely generated. In particular the group $\pi_1(X \setminus V)$ is finitely generated. Let us recall the following result of M. Hall (see [Hal], [Kur]):

Lemma 2.1. *Let G be a finitely generated group and let k be a natural number. Then there are only finite number of subgroups $H \subset G$ such that $[G : H] = k$.*

Every finite mapping $f : X \to Y$ with $D(f) = V$ and $\mu(f) = k$ induces a topological covering $f : X \setminus f^{-1}(D(f)) = P_f \to R = Y \setminus D(f)$. Take a point $a \in R$ and let $a_f \in f^{-1}(a)$. We have an induced homomorphism

$$f_* : \pi_1(P_f, a_f) \to \pi_1(R, a).$$

Denote $H_f = f_*(\pi_1(P_f, a_f))$ and $G = \pi_1(R, a)$. Hence $[G : H_f] = k$. By Lemma 2.1 there are only a finite number of subgroups $H_1, \ldots, H_r \subset G$ with index k. Choose finite regular mappings $f_i : X \to Y$ such that $H_{f_i} = H_i$ (of course only if such a mapping f_i does exist). We show that every finite regular mapping f such that $D(f) = V$ and $\mu(f) = k$ is equivalent to one of the mappings f_i.

Indeed, let $H_f = H_{f_i}$. We show that f is equivalent to f_i. Let us consider two coverings $f : (P_f, a_f) \to (R, a)$ and $f_i : (P_{f_i}, a_{f_i}) \to (R, a)$. Since $f_*(\pi_1(P_f, a_f)) = f_{i*}(\pi_1(P_{f_i}, a_{f_i}))$ we can lift the covering f to a homeomorphism $\phi : P_f \to P_{f_i}$ such that the following diagram commutes:



Note that $\phi : P_f \to P_{f_i}$ is a single-valued branch of the multi-valued holomorphic mapping $f_i^{-1} \circ f$. In particular ϕ is a holomorphic mapping.

Now we show that ϕ has a unique extension to a regular mapping on the whole of X. Let $x \in X$ and $y = f(x)$. Let U be an affine neighborhood of y in Y and take $X_1 = f^{-1}(U)$, $X_2 = f_i^{-1}(U)$. Since f, f_i are finite, the varieties X_1, X_2 are affine. Moreover, $\phi(X_1) \subset X_2$.

Since X_2 is affine, we have an embedding $X_2 \subset \mathbb{C}^N$ for N large enough. Consider the mapping $\phi' : P_f \cap X_1 \ni x \mapsto \phi(x) \in X_2 \subset \mathbb{C}^N$. Since the mappings f and f_i are proper the mapping ϕ' is locally bounded on X_1. Hence by the Riemann Extension Theorem, ϕ' can be extended to a holomorphic mapping $\Phi_{|X_1} : X_1 \to \mathbb{C}^N$. Of course we have $\Phi_{|X_1}(X_1) \subset X$. Gluing all possible mappings $\Phi_{|X_1}$ we get a global holomorphic extension $\Phi : X \to X$ of the mapping ϕ. Moreover we still have $f = f_i \circ \Phi$.

Finally, for fixed X_1 the graph of $\Phi_{|X_1}$ is a closed irreducible analytic n-dimensional subset of the n-dimensional affine variety $\Gamma = X_1 \times_Y X_2$ (the fiber product given by the mappings f and f_i). Consequently, the graph$(\Phi_{|X_1})$ has to coincide with some irreducible component of Γ, i.e., graph(Φ) is an algebraic subset of $X_1 \times X_2$. By the Serre Theorem about the algebraic graph (see e.g., [Łoj], p. 342) we know that $\Phi_{|X_1}$ is a regular mapping. Hence also Φ is regular. In a similar way the mapping Ψ determined by ϕ^{-1} is regular. It is easy to see that $\Psi \circ \Phi = \Phi \circ \Psi = \mathit{identity}$, hence Φ is a regular automorphism. Consequently, f is equivalent to f_i. □

Remark 2.2. It is easy to see from the proof that the number of non-equivalent finite regular mappings $f : X \to Y$ such that $D(f) = V$ and $\mu(f) = k$ is bounded by the number of subgroups of index k of the group $\pi_1(Y \setminus V)$.

Remark 2.3. Assume that X, Y are holomorphic and additionally the group $\pi_1(Y \setminus V)$ is finitely generated. Then the holomorphic counterpart of Theorem 1.3 is still true (this confirms Remark 1.4). The proof is the same as above, with one exception: we have to prove in a different way that the mapping ϕ has a holomorphic extension to the whole of X. To do it take a point $x \in f^{-1}(V)$ and let $y = f(x)$. The set $f_i^{-1}(y) = \{b_1, ..., b_s\}$ is finite. Take small open disjoint neighborhoods W_i of b_i, such that each W_i is biholomorphic to a small ball in $\mathbb{C}^n$ (here $n = \dim X$). We can choose an open neighborhood Q of y so small that $f_i^{-1}(Q) \subset \bigcup_{j=1}^{s} W_i$. Now take a small connected neighborhood P_x of x such that P_x is biholomorphic to a small ball in $\mathbb{C}^n$ and $f(P_x) \subset Q$. The set $P_x \setminus f^{-1}(V)$ is still connected and it is transformed by ϕ into one particular set W_{i_0}. Now we can use the Riemann Extension Theorem.

3. Proof of Corollaries 1.4-1.11

Proof of Corollary 1.5. Throughout this section, we keep notation of Section 2. Note that in this case $G = \mathbb{Z}$ and for every k there is only one subgroup of G with index k. In particular we can choose f_1 as

$$f_1(x_1, x_2, \ldots, x_n) = (x_1^k, x_2, \ldots, x_n).$$

Arguing as above we see that every proper polynomial (holomorphic) mapping $f : \mathbb{C}^n \to \mathbb{C}^n$ with $D(f) = V$ and $\mu(f) = k$ is equivalent to f_1. $\square$

Proof of Corollary 1.6. In this case again we have $G = \mathbb{Z}$. Let $f, g : X \to \mathbb{C}^n$ be proper mappings with $D(f) = D(g)$ and $\mu(f) = \mu(g) = k$. Since G has only one subgroup of index k, we see as before that f is equivalent to g. In the case of Stein manifolds use the fact, that $R := \mathbb{C}^n \setminus V$ is Stein, and since an analytic covering of a Stein space is again Stein, we also see that $P_f = X \setminus f^{-1}(V)$ and $P_g = X \setminus g^{-1}(V)$ are Stein. Then one follows the proof of Theorem 1.3.

Moreover, since the group H_f is normal in G, the covering $f : P_f \to R$ is regular. In particular, the covering group $G(P_f|R)$ acts transitively on P_f. Since we know all covers of the space $\mathbb{C}^n \setminus V = \mathbb{C}^* \times \mathbb{C}^{n-1}$ (it has the homotopy type of a circle), we see that this group is cyclic. Now take as σ the generator of the group $G(P_f|R)$. As before, σ is a single-valued branch of the multi-valued holomorphic function $f^{-1} \circ f$, hence it is holomorphic. Again we can extend σ to the whole of X. Moreover, if f is a regular mapping, then (as above) so is σ. $\square$

Proof of Corollary 1.7. Let X be an algebraic (resp. holomorphic) manifold. As above, the mapping f induces a topological regular covering $f : P_f = X \setminus f^{-1}(V) \to R = Y \setminus V$. For a point $a \in R$ let a_1, a_2 be two different points in the fiber $f^{-1}(a)$. Since f is a regular covering (the topological degree is two) there exists a covering homeomorphism $\phi : P_f \to P_f$ such that $\phi(a_1) = a_2$. As before, the mapping ϕ is regular (resp. holomorphic).

Let $x \in X$ be such that $y = f(x) \in V$, in particular $f^{-1}(y) = x$. Let $U \cong B(0, r)$ be a small neighborhood of x in X. Since $f \circ \phi = f$ and f is finite, there is a small neighborhood U' of x such that $\phi(U' \setminus f^{-1}(V)) \subset U$. The mapping $\phi : U' \setminus f^{-1}(V) \to U \cong B(0, r) \subset \mathbb{C}^n$ is bounded, hence by the Riemann Theorem it can be extended to a holomorphic mapping $\Phi : U' \to U$ and $\Phi(x) = x$ for $x \in f^{-1}(V)$. This implies $Fix(\Phi) = C$ (C is the set of critical points of f). $\square$

Proof of Corollary 1.8. In virtue of Remark 1.17 we have that there is a non-trivial automorphism $\Phi : X \to X$ such that $f \circ \Phi = f$. Take $G = \{id, \Phi\}$. Since the mapping f is G-equivariant, we have a well-defined morphism $f' : X/G \ni [x] \mapsto f(x) \in Z$. By the assumption, the mapping f' is a bijection. By the Zariski Main Theorem (resp. by its holomorphic analogon), we have that f' is an isomorphism. Hence the mapping $f = {f'}^{-1} \circ \pi$ is equivalent to the projection π. $\square$

Remark 3.1. We have used here the fact that every subgroup of G of index two is normal. Note that this purely algebraic property is one of the reasons why the case $\mu(f) = 2$ is so special.

Proof of Corollary 1.9. Note that $\mathbb{C}^n \setminus K_n \cong \prod_{i=1}^n \mathbb{C}^*$, hence $G = \mathbb{Z} \oplus \ldots \oplus \mathbb{Z}$ (n times). It is easy to see that every subgroup of index k in G, which comes from finite mapping

f has the form $H(d_1,\dots,d_n) := H_1 \oplus \dots \oplus H_n$, where $(\mathbb{Z} : H_i) = d_i$ and $\prod_{i=1}^n d_i = k$. Indeed, take $F_i = \{x \in \mathbb{C}^n : f_i = 0\}$ and $W_i = \{x \in \mathbb{C}^n : x_i = 0\}$. Consider groups $H_1(\mathbb{C}^n \setminus \bigcup_{i=1}^n W_i) \cong \pi_1(\mathbb{C}^n \setminus K_n) = \mathbb{Z}^n$ and $H_1(\mathbb{C}^n \setminus \bigcup_{i=1}^n F_i) \cong f_*(\pi_1(\mathbb{C}^n \setminus \bigcup_{i=1}^n F_i) \subset G = \mathbb{Z}^n$. This implies that $H_1(\mathbb{C}^n \setminus \bigcup_{i=1}^n F_i) = \mathbb{Z}^n$. By the Mayer-Vietoris sequence we have

$$H_1(\mathbb{C}^n \setminus \bigcup_{i=1}^n F_i) = \bigoplus_{i=1}^n H_1(\mathbb{C}^n \setminus F_i)$$

and

$$H_1(\mathbb{C}^n \setminus K_n) = \bigoplus_{i=1}^n H_1(\mathbb{C}^n \setminus W_i).$$

The induced mapping $H_1(f) : \bigoplus_{i=1}^n H_1(\mathbb{C}^n \setminus F_i) \to \bigoplus_{i=1}^n H_1(\mathbb{C}^n \setminus W_i)$ coincide with the mapping $H_1(f) : H_1(\mathbb{C}^n \setminus \bigcup_{i=1}^n F_i) \to H_1(\mathbb{C}^n \setminus K_n)$ and shows that indeed the group $H(d_1,...,d_n)$ has a desired form.

The mapping $f_{d_1,\dots,d_n} : \mathbb{C}^n \ni (x_1,\dots,x_n) \mapsto (x_1^{d_1},\dots,x_n^{d_n}) \in \mathbb{C}^n$, restricted to the set $\mathbb{C}^n \setminus K_n$, satisfies

$$f_{d_1,\dots,d_n *}(\pi_1(\mathbb{C}^n \setminus K_n, a_f)) = H(d_1,\dots,d_n).$$

This implies that every proper polynomial (resp. holomorphic) mapping with $D(f) = K_n$ and $\mu(f) = k$ has to be equivalent to one of the mappings $f_{d_1,\dots,d_n}$. However, we have to exclude mappings for which some of d_i is equal to one, because in this case $D(f)$ does not contain the hyperplane $x_i = 0$. □

Proof of Corollary 1.10. The mapping f induces a topological covering $f : P_f = \mathbb{C}^n \setminus f^{-1}(K_n) \to R = \mathbb{C}^n \setminus K_n$ of degree $\mu(f)$. Choose a point $a \in R$ and a point $a_f \in f^{-1}(a)$. Let $H(d_1,\dots,d_n)$ be the group $f_*(\pi_1(P_f, a_f)) \subset \pi_1(R, a)$.

As in the proof of Corollary 1.9 there is a mapping $f_{d_1,\dots,d_n} : \mathbb{C}^n \ni (x_1,\dots,x_n) \mapsto (x_1^{d_1},\dots,x_n^{d_n}) \in \mathbb{C}^n$ (and a point $b \in f^{-1}_{d_1,\dots,d_n}(a)$), which when restricted to the set $\mathbb{C}^n \setminus K_n$, satisfies $f_{d_1,\dots,d_n *}(\pi_1(\mathbb{C}^n \setminus K_n, b)) = H(d_1,\dots,d_n)$.

As before, we have a biholomorphic mapping $\phi : P_f \to P_{f_{d_1,\dots,d_n}}$ such that $f = f_{d_1,\dots,d_n} \circ \phi$. Since the mapping $f_{d_1,\dots,d_n}$ is proper, we can extend ϕ to a unique birational mapping $\Phi : \mathbb{C}^n \to \mathbb{C}^n$ and still have $f = f_{d_1,\dots,d_n} \circ \Phi$. However, the mapping f is quasi-finite, hence so is Φ. Now by the Zariski Main Theorem the mapping Φ is a regular automorphism. □

Proof of Corollary 1.11. In dimension one the hypersurface $X \subset \mathbb{C}$ which contains one or two points has only one embedding in $\mathbb{C}$ (up to equivalence). In dimension two by the famous Abhyankar-Moh-Suzuki Theorem the line $\mathbb{A}^1(\mathbb{C})$ has only one embedding in $\mathbb{C}^2$. Finally the $n-$cross $K_n = \{x \in \mathbb{C}^n : \prod_{i=1}^n x_i = 0\}$ has only one embedding into $\mathbb{C}^n$ (see [Jel]). By the remarks above, we can choose the polynomial automorphism $\Psi : \mathbb{C}^n \to \mathbb{C}^n$ such that $D(\Psi \circ f)$ is equal to:

a) the point 0 or the set $\{0, 1\}$ respectively,

b) the line $\{x = 0\} \subset \mathbb{C}^2$,

c) the n-cross $K_n \subset \mathbb{C}^n$.

Now it is enough to apply Corollary 1.5 or Corollary 1.9.□

4. Proofs of Theorem 1.12 and Corollaries 1.13-1.14

Proof of Theorem 1.12. Let $B(0,\rho)$ be a small ball around the origin. It is well known that the fundamental group of the space $U_\rho = B(0,\rho) \setminus V$ does not depend on sufficiently small ρ, and this group is finitely generated. Moreover, it is easy to see that for small $\rho_1 < \rho$ the space U_{ρ_1} is a deformation retract of U_ρ.

Every proper mapping $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ with $D(f) = (V, 0)$ and $\mu(f) = k$ induces a topological covering $f : f^{-1}(U_\rho) = P_{f,\rho} \to U_\rho$. Take a point $a \in U_\rho$ and let $a_f \in f^{-1}(a)$. We have an induced homomorphism

$$f_* : \pi_1(P_{f,\rho}, a_f) \to \pi_1(U_\rho, a).$$

Denote $H_{f,\rho} = f_*(\pi_1(P_{f,\rho}, a_f))$ and $G = \pi_1(U_\rho, a)$. Note that for small $\rho_1 < \rho$ the space P_{f,ρ_1} is a deformation retract of $P_{f,\rho}$. In particular the group $H_{f,\rho}$ does not depend on ρ; we will denote it by H_f. We also write P_f instead of $P_{f,\rho}$, and U instead of U_ρ.

Hence $[G : H_f] = k$. By Lemma 2.1 there are only a finite number subgroups $H_1, \ldots, H_r \subset G$ with index k. Choose proper holomorphic mappings $f_i : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ such that $H_{f_i} = H_i$ (of course only if such a f_i exists). We show that every proper holomorphic mapping $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ such that $D(f) = (V, 0)$ and $\mu(f) = k$ is equivalent to one of the f_i.

Indeed, let $H_f = H_{f_i}$. We show that f is equivalent to f_i. Let us consider two coverings $f : (P_f, a_f) \to (U, a)$ and $f_i : (P_{f_i}, a_{f_i}) \to (U, a)$. Since $f_*(\pi_1(P_f, a_f)) = f_{i*}(\pi_1(P_{f_i}, a_{f_i}))$ we can lift the covering f to a homeomorphism $\phi : P_f \to P_{f_i}$ such that the following diagram commutes:



Note that $\phi : P_f \to P_{f_i}$ is a single-valued branch of the multi-valued holomorphic mapping $f_i^{-1} \circ f$. In particular ϕ is a holomorphic mapping. Further, the mapping ϕ is bounded on P_f. Hence by the Riemann Extension Theorem ϕ can be extended to a holomorphic mapping $\Phi : f^{-1}(B(0,\rho)) \to f_i^{-1}(B(0,\rho))$. Moreover we still have $f = f_i \circ \Phi$.

In a similar way the mapping $\Psi : f_i^{-1}(B(0,\rho)) \to f^{-1}(B(0,\rho))$ determined by ϕ^{-1} is holomorphic. It is easy to see that $\Psi = \Phi^{-1}$, hence Φ is a biholomorphism. Consequently, f is equivalent to f_i. □

Proof of Corollary 1.13. Let $U_\rho = B(0,\rho) \setminus V$. It is easy to see that U_ρ is a deformation retract of $\mathbb{C}^n \setminus V$. In particular $G = \mathbb{Z}$. Now the rest of the proof of the first part of Corollary 1.13 is the same as in the proof of Corollary 1.5 so we skip it.

Now assume that $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ is a proper holomorphic mapping with a smooth discriminant $D(f)$. Let $\Psi : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ be a biholomorphism such that $\Psi(D(f)) = (V, 0)$, where $V = \{x : x_1 = 0\}$. Hence $D(\Psi \circ f) = (V, 0)$ and by the first part $\Psi \circ f$ is equivalent to g.

Proof of Corollary 1.14. Again U_ρ is a deformation retract of $\mathbb{C}^n \setminus K_n$. In particular $G = \mathbb{Z} \oplus \ldots \oplus \mathbb{Z}$ (n times). Now the rest of the proof of the first part of Corollary 1.14 is the same as in the proof of Corollary 1.9 so we skip it.

Assume now that $D(f)$ is biholomorphic to K_n, and let $\sigma : (K_n, 0) \to (D(f), 0)$ be this isomorphism. Let $\Lambda_i = \sigma(\{x : x_i = 0\})$ and let $h_i = 0$ be a local reduced equation of Λ_i. Take $\Psi = (h_1, \ldots, h_n)$. Then Ψ is a local biholomorphism (all Λ_i intersect transversally at 0 !) and it is easy to see that $D(\Psi \circ f) = (K_n, 0)$. Now we can apply the first part of the Corollary. □

5. Proofs of Theorems 1.15 and 1.16

Proof of Theorem 1.15. Let $f : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ be a proper holomorphic mapping of degree two. Let $B(0, \rho)$ be a small ball around the origin and $U_\rho = B(0, \rho) \setminus V$. Take a point $a \in U_\rho$ and let $b_1, b_2 \in f^{-1}(a)$. We have the induced homomorphisms

$$f_* : \pi_1(P_f, b_i) \to \pi_1(U_\rho, a), \ i = 1, 2.$$

Denote $H_i = f_*(\pi_1(P_f, b_i))$ for $i = 1, 2$ and denote as above $G = \pi_1(U_\rho, a)$.

Since $[G : H_i] = 2$ we see that each H_i is a are normal subgroup of G. From topology we know that H_1 is conjugate to H_2, and consequently $H_1 = H_2$. Let us consider two coverings $f : (P_f, b_1) \to (U, a)$ and $f : (P_f, b_2) \to (U, a)$. Since $f_*(\pi_1(P_f, b_1)) = f_*(\pi_1(P_f, b_2))$ we can lift the former covering to a homeomorphism $\phi : (P_f, b_1) \to (P_f, b_2)$ such that the following diagram commutes:



Put $R = f^{-1}(B(0, \rho))$. As above, ϕ can be extended to a biholomorphism $\Phi : R \to R$ of order two with $f = f \circ \Phi$. Since Φ permutes the fibers of f, we see that all critical points of f are fixed points of Φ. In particular $\Phi(0) = 0$. Now by the Cartan Theorem (see [Car]) we know that for ρ small enough, there is a biholomorphism $\Sigma : (R, 0) \to (R, 0)$ such that $\Sigma^{-1} \circ \Phi \circ \Sigma := L$ is a linear automorphism. Now we need the following:

Lemma 5.1. *Let $L : \mathbb{C}^n \to \mathbb{C}^n$ be a linear mapping of order two and assume that the set of fixed points of L is a hyperplane W. Then in some coordinates,*

$$L(x_1, x_2, \ldots, x_n) = (-x_1, x_2, \ldots, x_n).$$

Proof. Take a basis $e_1, \ldots, e_n$ in $\mathbb{C}^n$ such that $e_2, \ldots, e_n$ span the hyperplane W. Hence $L(e_i) = e_i$ for $i > 1$ and $L(e_1) = \sum_{i=1}^n a_i e_i$. In particular $L(x_1e_1 + \ldots + x_ne_n) = (a_1x_1)e_1 + (x_2 + a_2x_1)e_2 + \ldots + (x_n + a_nx_1)e_n$. Since L has finite order, we have $a_i = 0$ for $i > 1$. In particular $L(x_1, \ldots, x_n) = (a_1x_1, x_2, \ldots, x_n)$. However $\det L = \pm 1$, i.e., $a_1 = \pm 1$. Of course $a_1 \neq 1$. □

Let L be a linear automorphism as in Lemma 5.1 and put $G = (id, L)$. The group G acts on $\mathbb{C}^n$ and we have $\mathbb{C}[x_1, \ldots, x_n]^G = \mathbb{C}[x_1^2, x_2, \ldots, x_n]$. In particular

$$\mathbb{C}^n/G = Spec(\mathbb{C}[x_1, \ldots, x_n]^G) = Spec(\mathbb{C}[x_1^2, x_2, \ldots, x_n]) = \mathbb{C}^n.$$

Under this identification the canonical mapping $\pi : \mathbb{C}^n \to \mathbb{C}^n/G$ coincides with the mapping $\mathbb{C}^n \ni (x_1, \ldots, x_n) \mapsto (x_1^2, x_2, \ldots, x_n) \in \mathbb{C}^n$.

Take $f' = f \circ \Sigma$. Hence f' is equivalent to f and $f' \circ L = f'$. The mapping f' induces a mapping $\tilde{f} : R/G = \pi(R) \to B(0, \rho)$, where $f' = \tilde{f} \circ \pi$. Since $\tilde{f}$ has degree one it is a biholomorphism. However $\tilde{f}^{-1} \circ f' = \pi$, so the mapping f' (hence also f) is weakly equivalent to

$$\pi : (\mathbb{C}^n, 0) \ni (x_1, \ldots, x_n) \mapsto (x_1^2, x_2, \ldots, x_n) \in (\mathbb{C}^n, 0). \square$$

Proof of Theorem 1.16. By Corollary 1.7 there exists a non-trivial automorphism $\Phi : \mathbb{C}^n \to \mathbb{C}^n$ such that $f \circ \Phi = f$. In particular the set of fixed points of Φ coincide with the set of critical points of f. If the Linearization Conjecture is true, then $\Sigma^{-1} \circ \Phi \circ \Sigma = L \in GL(n)$ for some automorphism $\Sigma \in Aut(\mathbb{C}^n)$. As above we can assume that $L(x_1, x_2, \ldots, x_n) = (-x_1, x_2, \ldots, x_n)$. Put $f := f \circ \Sigma$. Now we have $f \circ L = f$.

If $G = (id, L)$, then we have $\mathbb{C}^n/G = \mathbb{C}^n$ and the canonical mapping $\pi : \mathbb{C}^n \to \mathbb{C}^n/G$ coincides with the mapping $\mathbb{C}^n \ni (x_1, \ldots, x_n) \mapsto (x_1^2, x_2, \ldots, x_n) \in \mathbb{C}^n$. Since f is G-invariant, the mapping

$$\tilde{f} = f \circ \pi^{-1} : \mathbb{C}^n/G \ni (x_1, \ldots, x_n) \mapsto f(\sqrt{x_1}, x_2, \ldots, x_n) \in \mathbb{C}^n$$

is well defined and polynomial. It is easy to check that $\tilde{f}$ has degree one, i.e., it is an automorphism. Finally $\tilde{f}^{-1} \circ f = \pi$. □

In particular for $n = 2$ we get another proof of the Lamy Theorem (Theorem 1.1).

References

[B-C] C. Bisi, F. Polizzi, On Proper Polynomial Maps of $\mathbb{C}^2$, *Journal of Geometric Analysis 20*, (2010), 72-89.

[B-C-1] C. Bisi, F. Polizzi, Proper polynomial self-maps of the affine space: state of the art and new results, *Contemporary Mathematics 553* (2011), 15-25.

[Car] H. Cartan, Sur le groupes de transformations analytiques, *Actualities Scientifiques et Industrielles, 198*, Paris 1935.

[Hal] M. Hall, A topology for free groups and related topics, *Annals Math. 52* (1950), 127-139.

[Jel] Z. Jelonek, A hypersurface which has the Abhyankar-Moh Property, *Math. Ann. 308*, 73-84, (1997).

[J-K] Z. Jelonek, K. Kurdyka, Quantitative Generalized Bertini-Sard Theorem for smooth affine varieties, *Discrete and Computational Geometry 34*, 659-678, (2005),

[Kam] T. Kambayashi, Automorphism group of a polynomial ring and algebraic group action on an affine space, *J. Algebra 60*, 439-451, (1979).

[Kur] A. G. Kurosh, *Group theory*, Chelsea Publishing Company, 1960.

[Lam] S. Lamy, Sur la structure du groupe d´ automorphismes de certaines surfaces affines, *Publ. Mat. 49* (2005), 3-20.

[Łoj] S. Łojasiewicz, *Introduction to complex analytic geometry*, PWN, Warszawa, 1988.

(Z. Jelonek) Instytut Matematyczny, Polska Akademia Nauk, Śniadeckich 8, 00-956 Warszawa, Poland

E-mail address: najelone@cyf-kr.edu.pl